

# Heliotropium indicum L: An Ethnomedicinally Important plant of India Rupam Mandal

Keywords: Ethnomedicinally Important, *Heliotropium indicum*, phytopharmacological activities, Medicinal Value.

#### Abstract:

This study attempted to summarise the existing data on the ethnomedicinal and phytopharmacological activities of *Heliotropium indicum* L., and it did so by using database reports as its primary source of information. In order to accomplish this goal, a search of the most recent relevant literature was carried out. Based on the data, it appears that the plant possesses a significant quantity of essential phytochemicals. In the plant family known as the Boraginaceae, *Heliotropium* is a sizable genus that spans both the tropical and temperate regions of the world. Because most of the alkaloids that can be isolated from *Heliotropium indicum* are toxic, this plant's use in treating internal conditions is not suggested. In order to provide light on potential directions for future research, the purpose of this review is to summarise the phytochemistry and pharmacological activity of the many species of *Heliotropium*.

### **Introduction:**

New medicinal agents are frequently discovered using natural products. Traditional medicine has been the most logical and cost-effective form of care. From ancient time plant have been used as a source of therapy (Maiti et al., 2013; Sanyal, 2016). The medicinal plant plays a vital role in discovering new therapeutic agents (Banerjee et al., 2014; Bhattacharjee and Manna, 2016). Various phytochemicals like alkaloids, flavonoids, tannins, and phenols contribute to the medicinal property of medicinal plants. The medicinal use of herbs is deeply rooted in Indian culture (Maiti et al., 2010; Sarkar et al., 2016; Sarkar, 2017; Bhattacharjee, 2021). Traditional systems of medicine, along with homoeopathy and folklore medicine, both of which are practised in India, continue to play an essential part in the country's overall healthcare system for the general people (Sanyal et al., 2018; Erfani, 2021; Kundu, 2022; Kar et al., 2022).

*Heliotropium indicum* is locally known as Hatisur and gets its name from the Greek word "helios" meaning sun and "tropein" meaning to turn indicating that its flower turn towards the sun. This plant is distributed throughout Bangladesh, Sri Lanka, Thiland, Nepal, India, tropical

#### Rupam Mandal

Department of Botany, Sree Chaitanya College, Habra, West Bengal, India. E-mail: sanphuip@gmail.com Orcid iD: https://orcid.org/0000-0002-0037-3726 Author's E-mail: sanphuip@gmail.com

© International Academic Publishing House, 2022 Bhanumati Sarkar (eds.), The Basic Handbook of Indian Ethnobotany and Traditional Medicine, Vol. 1 ISBN: 978-81-957954-1-3 Published online: 23rd August, 2022 Asia, and some parts of Africa. This plant is traditionally used against many pathological diseases (Sarkar, 2016; Chakraborty et al., 2019). In this chapter, the ethnomedicinal, phytochemical, and pharmacological profiles of *Heliotropium indicum* are intended to be reflected in the current context.

# **Plant Taxonomy:**

Heliotropium indicum belongs to-Domain: Eukaryota Kingdom: Plantae Phylum: Spermatophyta Subphylum: Angiospermae Class: Dicotyledonae Order: Boraginales Family: Boraginaceae Genus: Heliotropium Species: indicum

# **Plant Morphology:**

*Heliotropium indicum* is an annual, perennial herb reaching a height of 20 to 50 cm (Dash and Abdullah, 2012). The stem is widely distributed, branched or unbranched. Long and heavily branched tap-roots make up the root system. The leaves are oval or ovate-oblong, simple alternate or sub-alternate, and have an undulate border. The undersides of the leaves are visibly covered in hair and nerve, which are found on both sides of the serrulate or cordate leaves. Heliotropium indica blooms all year long. Within the cymose, the flower grows apically. The fruit is referred to as a nutlet. They are 4-6 mm long, 2-4 lobed, and indehiscent. There are two cells and a beak on each nutlet. At the base of the flower, mature nutlets are visible. The flower is sessile, two-ranked pentamerous, and white or violet in colour. The sepals are five, evenly spaced with the hair outside, deep green in colour, irregular, and between three and five millimetres long.

# **Traditional and folk values:**

Different traditional and folkloric medical systems have employed *Heliotropium indicum* to treat various ailments. *Heliotropium indicum* is a medication used to treat neurological problems, poison ivy, and skin conditions in Tamil Nadu, India. According to legend, the Malasar tribe used coconut oil and leaf juice boiled together to treat dandruff (Rahman et al., 2011). The root juice of *Heliotropium indicum* is used by some tribes in Assam, India, to treat ophthalmia.

When consumed orally, the dried flower is said to sterilize females permanently. Thailand is home to this practice. *Heliotropium indicum* is thought to be beneficial in treating dermatitis, stomach pain, and malaria in various African nations (Rahman et al., 2011). The flower is also infused and used to treat menorrhagia in Jamaica.

Several ailments have been treated using the entire plant in various traditional systems. In Bangladesh, the extract of *Heliotropium indicum* is used to cure chicken pox, allergies, knee swelling, and poisoning (Nawaz, 2009).

In the Philippines, dried root decoction is used to promote menstruation, flower infusion is used to cure kidney stones, and seeds are used to heal wounds, cholera, and malaria (Wiart, 2006). In Malaysia, the whole plant decoction is used to treat ringworm infection, whereas it is used to treat gonorrhoea in Burma (Wiart, 2006). Table 1 includes a list of the regional names and traditional uses of *Heliotropium indicum* in several nations.

Country	Local name	Traditional use for	Plant parts used	Reference
Bangladesh	Hatisur	Antidote to poisoning	Leaves and stem	Nawaz, 2009
		Swelling of Knee, joint pain	Root	Nawaz, 2009
		Chicken pox, Allergy	leaves	Shahnaj et al., 2015
		Blood purification	Root	Akhter et al., 2021
Benin	Koklosouden	Psychosis	Leaf and root	Adjanohoun, 2011
		Leucorrhoea	Whole plant	Adjanohoun, 2011
Ghana	Komfemtikoro	Eye infection	leaves	Komlaga, 2017
India	Nakipoo	Snake and scorpion bite	leaves	Alagesaboopathi, 2009
	Indian heliotrope & Hatisur	Wound and skin infection	Whole plant	Muthu, 2006
		ophthalmia	root	Das et al., 2008
Indonasia	Bandotanlombok	Herpes & rheumatism	leaves	Togola, 2005
Jamaica	Turnsoles	Menorrhagia	flower	Asprey and
				Thornton, 1955
		Fever, ulcer, sore	Whole plant	Asprey and
		throat, Induced		Thornton, 1955
		abortion, rectal sores		
Mauritius	Herbepapillon	Renal colic	leaves	Suroowan and
				Mahomoodally,
				2019
Mali	Bambara	Nausea & vomit-ing,	leaves	Nordeng et al.,
		high blood pressure		2005

### Table 1. The use of H.indicum in traditional medicine

Heliotropium indicum L: An Ethnomedicinally Important plant of India

Nigeria	Akuko	Hepatitis and fever	leaves	Adjanohoun, 2014
		Gonorrhea	leaves	Ainslie et al., 1981
Philippines	Buntot-leon	Diuretic & kidney	Whole plant	Quisumbing, 1951
		stone		
Senegal	Mandingbambaranagik	Dermatitis	leaves	Kerharo and Adam,
	u			1974
Taiwan	Gou Wei chungtsan	Hepatitis	Leaves and	Lin and Kan,1990
			root	
West Indies		Head lice	Whole plant	Ayensu, 1981

# **Phytochemical Constituent**

As *Heliotropium indicum* has been widely used in traditional and folk medicine since ancient times, researchers have made many efforts to identify the different phytochemicals and pharmacologically active compounds that contribute to the wide use of this plant in herbal medicine. Reports suggest that the areal part of *Heliotropium indicum* contains different alkaloids like helindicine, echinitine, europine, indicinine, heliotrine-Noxide. The roots contains a high level of estradiol. *Helindicine* is a new pyrrolizidine alkaloid that has been isolated from the roots of Heliotropium indicum (Souza et al., 2005). The plant is reported to contain triterpenes like lupeol, rapone, rapanone, amines like putrescine, spermidine, spermine and sterols like astradiol, campesterol etc. (Wiart, 2006 ; Coe and Anderson,1996; Lin and Kan,1990). *Heliotropium indicum* also yields an essential oil that consists mainly of phytol, 1-dodecanol and beta-linalool. The different chemical compounds isolated from the plant are listed in **table 2**.

Plant part	Phytochemicals	References	
Seed	Alkaloids:	Williaman and Schubert,	
	Cynoglossine, Europine N-oxide, Heleurine N-	1962	
	oxide, Heliotridine N-oxide, Heliotrine N-oxide		
Aerial part	Alkaloids:	Duke, 1994; Coe and	
	Echinatine, Heleurine, Heliotrine, Heliotridine,	Anderson,1996; Pandey et	
	Indicine Aerial, Indicine N-oxide, Lasiocarpine,	al.,1996	
Leaves	Alkaloid:	Birecka et al., 1984	
	Trachelanth-amidine		
	Amines:		
	Putrescine, Spermidine, Spermine		
Whole plant	Sterols:	Andhiwal, 2013; Machan et	
	β-Sitosterol, Estradiol, Chalinasterol,	al., 2005	
	Campesterol ,Hexacosane-1-ol, Stigmasterol		
	Volatile oils:1-Dodecanol, $\beta$ -Linalool, Phytol		

 Table 2. Different Phytochemicals present Heliotropium indicum

#### **Medicinal Importance:**

Based on the data available on the presence of various phytochemicals in the different solvent extracts of Heliotropium indicum and its enormous use in folk medicine, many investigations have been conducted to look into the medicinal importance of the plant. The results have shown diverse biological effects described in this section.

#### **Anti-microbial Activity:**

The alcoholic extracts of the whole plant exhibited a dose-dependent anti-bacterial property. The extracts were effective against Bacillus subtilis, Bacillus pumilus, Staphylococcus aureus, Pseudomonous aeruginosa, Escherichia coli, Proteus vulgaris (Osungunna and Adedeji, 2016; Rao et al., 2002). In another study, the petroleum ether, chloroform, aqueous and methanolic leaf extract of Heliotropiumindicumhas shown antibacterial effects against Bacillus subtilis, Escherichia coli, Pseudomonous aeruginosa and Staphylococcus aureus. (Oluwatoyin et al., 2011; Dash and Murthy, 2011). It has also been reported that the methanolic and aqueous leaf extracts mixed with simple ointment give promising results in *Staphylococcus aureus* induced wound infection model (Dash and Murthy, 2011). The volatile oils isolated from the aerial parts of the plant showed antituberculosis activity against Mycobacterum tuberculosis (Machan et al., 2005). The alcoholic extracts also possessed anti-fungal activity as tested against Aspergillus niger, Aspergillus wentii, Rhizopus oryzae, Candida albicans (Osungunna and Adedeji, 2016; Rao et al., 2002). However, the anti-bacterial and antifungal effect of Heliotropium indicum high extract concentration is required. Therefore, the isolation and characterization of the active components and formulation based on those active components would increase the efficiency of this plant's antimicrobial properties.

#### **Anti-inflammatory Activity:**

The leaf extract of *Heliotropium indicum* has exhibited anti-inflammatory activity carrageenan-induced hind paw oedema and cotton pellet granulosa model of inflammation. The methanolic root extract of *Heliotropium indicum* showed a significant anti-inflammatory effect, with a 49% reduction in paw edema and a 55% reduction in granuloma formation (Srinivas et al., 2000). The ethanolic and petroleum ether extracts of *Heliotropium indicum* could successfully reduce the inflammation in egg white and induce acute paw oedema in a rat model (Shalini et al., 2010). Aqueous whole plant extract of *Heliotropium indicum* has also been reported to show anti-inflammatory effects in lipopolysaccharide-induced rabbit models. The extracts reduce inflammation and inflammatory cell infiltration (Kyeiet et al., 2016). It is noted that an oral pharmaceutical product obtained from *Heliotropium indicum* is used against inflammation, particularly against inflammatory diseases of the intestine (Pianowski et al., 2011).

#### Wound healing capacity:

Different *Heliotropium indicum* extracts have been shown to have wound-healing properties. In a rat model, the alcoholic extract of *Heliotropium indicum* demonstrated wound-healing efficacy. Topically, rats who received an extract demonstrated full wound healing (Reddy et al., 2002). The alkaloids extracted from n-butanol crude extract display remarkable wound healing activity in human lung cell line H292.

### **Anticancer activity:**

*Heliotropium indicum* possesses anticancer properties, as is evident from a study on HeLa cells using stem and leaf methanolic extract. Within 48 hours of treatment, both extracts demonstrated anti-proliferative effects (Sivajothi et al., 2015). On the SKBR3 human breast cancer cell line, an ethanolic extract of *Heliotropium indicum* demonstrated an anti-proliferative activity (Goyal and Sharma, 2014). The brine shrimp nauplii were cytotoxic to the *Heliotropium indicum* methanolic root extract, with an LC50 value ranging from 2.57 to 31.44 ug/ml (Rahman et al., 2011). Interestingly, the main pyrrolizidine alkaloid derived from the plant, indicine N-oxide, has entered phase-I clinical trials in patients with advanced cancer (Ohnuma et al., 1982).

### **Antifertility activity:**

The ethnomedicinal use of *Heliotropium indicum* as an antifertility agent is supported by the plant's substantial antifertility and abortifacient action in the petroleum ether extract (Andhiwal et al., 2013). A study discovered that the n-hexane and benzene fractions of *Heliotropium indicum*'s ethanol extracts displayed antifertility effects in rat anti-implantation and abortifacient models (Savadi et al., 2009). The unfavourable impact of *Heliotropium indicum* extracts on sperm mortality was also demonstrated in the in-vitro study.

#### **Anticataract effect:**

Rats exhibit anticataract behaviour in response to the ethanolic leaf extract. According to the study, rats given extract treatment had much more water, soluble protein, and glutathione in their lenses (Veda et al., 2016). The aqueous extract of the entire plant *Heliotropium indicum* greatly reduced the occurrence of selenite-induced cataract (Kyei et al., 2015).

#### Anti helmintic effect:

On Indian adult earthworms, the leaf extracts of *Heliotropium indicum* in both methanol and water demonstrated an anti-helminthic activity. The effect of the methanolic extract was comparable to that of the reference medication, mebendazole. On the other hand, the aqueous extract had a significantly smaller and less potent effect on earthworms (Mahato et al., 2014).

#### **Conclusion:**

The conventional medical system makes extensive use of *Heliotropium indicum*. Numerous compounds with promising biological action may be obtained from the plant. Modern society is concerned about the overuse of medicinal plants for pharmacological purposes. According to reports, approximately 15000 plant species may go extinct as a result of overharvesting and habitat damage brought on by human activity. This should be considered while employing plants with medicinal value.

The use of herbal medicine has gained popularity recently, although there are still some questions about its effectiveness and safety. According to the article, *Heliotropium indicum* is used in several nations, but additional research is needed to identify the phytochemicals and therapeutic properties. The use of this plant in clinical practice is constrained because there have been relatively few investigations on its isolated ingredient, despite the fact that it can significantly progress medical treatment. Animal models for the toxicity of the extracts show contributions from heliotrine, lasiocarpine, and retrogressive presence. As a result, it is crucial to solve this issue. These problems can be solved with enough research, opening the door for this plant as a secure, efficient, and reasonably priced healthcare option.

### **References:**

- Adjanohoun, E. (2014) Le processus de r'enovation de la pharmacop'eeafricaine, *Bulletin de la Soci'et'eBotanique de France Actualit'es Botaniques*. 136(3-4): 35–39.
- Ainslie, J. R. (1981) Fe List of Plants Used in Native Medicine in Nigeria, Imp, Forest Inst Oxford Inst Paper. vol. 7.
- Akhter, J., Khatun, R. and Akter. S. (2021) Ethnomedicinal practices in Natore district, Bangladesh, *World Journal of Pharmacy and Pharmaceutical Sciences*, 5(8): 212–222
- Alagesaboopathi, C. (2009) Ethnomedicinal plants and their utilization by villagers in Kumaragiri hills of Salem district of Tamilnadu, India. *African Journal of Traditional, Complementary and Alternative Medicines*. 6 (3): 222–227.
- Andhiwal, C. K., Has, C., and Varshney, R. P., (2013) Chemical and pharmacological studies of Heliotropiumindicum. *Indian Drugs*. 22(11): 567–569.
- Asprey, G. F., and ornton, P. (1955) Medicinal plants of Jamaica. Parts III. *West Indian Medical Journal*. 4(4) : 69–82.
- Ayensu, E. S. (1981) *Medicinal Plants of the West Indies*, Reference Publications, Inc, Algonac, MI, USA
- Banerjee, J., Biswas, S., Madhu, N. R., Karmakar, S. Re., & Biswas, S. J. (2014). A better understanding of pharmacological activities and uses of phytochemicals of *Lycopodium clavatum*: A review. *Journal of Pharmacognosy and Phytochemistry*. 3 (1): 207-210.
- Bhattacharjee, P. (2021). Some medicinal plants with anti-fertility potential used by the tribal people of the District Cooch Behar, West Bengal, India. *International Journal of Experimental Research and Review*. 24: 30-39. doi: https://doi.org/10.52756/ijerr.2016.v03.006
- Bhattacharjee, P., & Manna, C. K. (2016). Potential plants as nervine for Unani system of medicine from the Coochbehar district, West Bengal, India. *International Journal of Experimental Research and Review*. 5: 19-24.
- Birecka, H., DiNolfo, T. E., Martin, W. B., and Frohlich, M. W. (1984) Polyamines and leaf senescence in pyrrolizidine alkaloidbearingHeliotropiumplants. *Phytochemistry*. 23(5): 991–997

Chakraborty, D., Das, D., Samal, A. C., and Santra, S. C. (2019). Prevalence and

Ecotoxicological significance of heavy metals in sediments of lower stretches of the Hooghly estuary, India. *International Journal of Experimental Research and Review*. 19: 1-17. doi: https://doi.org/10.52756/ijerr.2019.v19.001

- Coe F. G., and Anderson, G. J. (1996) Ethnobotany of the gar'ıfuna of Eastern Nicaragua, *Economic Botany*. 50 (1) : 71–107
- Das, A. K., Dutta, B. K., and Sharma, G. D. (2008) Medicinal plants used by different tribes of Cachar district, Assam. *Indian Journal of Traditional Knowledge*. 7 (3): 446–454.
- Dash, G. K., and Abdullah, M. S. (2012) A review on Heliotropiumindicum L. (Boraginaceae). International Journal of Pharmaceutical Sciences and Research. 4(4): 1253
- Duke, J. A. (1994) Amazonia Ethnobotanical Dictionary, CRC Press, Boca Raton, FL, USA.
- Erfani, H. (2021). The practical and potential importance of herbs such as ginger in Chemical Environmental Science. *International Journal of Experimental Research and Review*. 24:24-29. doi: https://doi.org/10.52756/ijerr.2021.v24.003
- Goyal, N., and Sharma, S. (2014) Bioactive phytoconstituents and plant extracts from genus *Heliotropium*. *InternationalJournal of Green Pharmacy*. 8(4): 217–22
- Kar, D., Ghosh, P., Suresh, P., Chandra, S., and Paul, D. (2022). Review on Phyto-chemistry & pharmacological activity of Melia azedarach. *International Journal of Experimental Research and Review*. 28: 38-46. https://doi.org/10.52756/ijerr.2022.v28.006
- Kerharo, J., and Adam, J. G. (1974) La pharmacopees ´ en´egalaise ´ traditionnelle: plantesm´edicinalesettoxiques, Editions VigotFr`eres, vol. 1011.
- Komlaga, G., Agyare, C., Dickson R. A., Annan, K, Loiseau, P.M., and Champy, P. (2015) Medicinal plants and finished marketed herbal products used in the treatment of malaria in the Ashanti region, Ghana. *Journal of Ethnopharmacology*. 172: 333–346.
- Kundu, K. (2022). Management of root-knot nematodes, Meloidogyne incognita in Okra using wheat flour as bionematocides. *International Journal of Experimental Research and Review*. 28: 8-14. doi: https://doi.org/10.52756/ijerr.2022.v28.002
- Kyei, S., Koffuor, G. A., Ramkissoon, P., Afari, C., and Asiamah, E. A. (2015) claim of anticataract potential of *Heliotropiumindicum*: a myth or reality? *Ophthalmology and Ferapy*. 4(2): 115–128.
- Kyei, S., Koffuor, G. A., Ramkissoon, P., Ameyaw, E. O., and Asiamah, E. A. (2016) Antiinflammatory effect of *Heliotropiumindicum* Linn on lipopolysaccharide-induced uveitis in New Zealand white rabbits. *International Journal of Ophthalmology*. 9(4): 528–535
- Lin, C. C., and Kan, W. S. (1990) Medicinal plants used for the treatment of hepatitis in Taiwan. *American Journal of Chinese Medicine*. 181(1-2): 35–43.
- Machan, T., Korth, J., Liawruangrath, B., Liaewruangrath, S., and Pyne, S. (2005) Composition and antituberculosis activity of the volatile oil of Heliotropiumindicum Linn. growing in Phitsanulok, ailand. *Flavour and Fragrance Journal*. 21(2): 265–267.
- Mahato, K., Kakoti, B. B., Borah, S., and Kumar, M. (2014) Evaluation of *in-vitro* anthelmintic activity of *Heliotropiumindicum*Linn. leaves in Indian adult earthworm. *AsianPacific Journal of Tropical Disease*. 4: S259–S262.

The Basic Handbook of Indian Ethnobotany and Traditional Medicine, Vol. 1

- Maiti, A., Madhu, N. R. and Manna, C. K. (2013). Natural products traditionally used by the tribal people of the Purulia district, West Bengal, India for the abortifacient purpose. *International Journal of Traditional Medicine* (TANG). 3(2): e14. doi: http://dx.doi.org/10.5667/tang.2012.0045
- Maiti, A., Madhu, N.R. and Manna, C. K. (2010). Ethnomedicine used by the tribal people of the district Purulia, W. B., India in controlling fertility : and experimental study. Pharmacologyonline. 1: 783-802.
- Muthu, C., Ayyanar, M., Raja, N., and Ignacimuthu S. (2006) Medicinal plants used by traditional healers in Kancheepuram district of Tamil Nadu, India. *Journal of Ethnobiology and Ethnomedicine*. 2: 43.
- Nawaz, A. H., Hossain, M., Karim, M., Khan, M., Jahan, R., and Rahmatullah, M. (2009) An ethnobotanical survey of Rajshahi district in Rajshahi division, Bangladesh, American-Eurasian Journal of Sustainable Agriculture. 3(2): 143–150.
- Ohnuma, T., Sridhar, K. S., Ratner, L. H., and Holland, J. F., (1982) Phase I study of indicine *N*-oxide in patients with advanced cancer. *Cancer Treatment Reviews*. 66(7): 1509–1515.
- Oluwatoyin, S., Ndukwe, G. I., and Joseph, A. (2011) Phytochemical and antimicrobial studies on the aerial parts of Heliotropiumindicum Linn. *Annals of Biological Research*. 2 (2): 129–136.
- Osungunna, M. O., and Adedeji, K. A. (2016) Phytochemical and antimicrobial screening of methanol extract of Heliotropiumindicum leaf. *Journal of Microbiology and Antimicrobials*. 3 (8): 213–216.
- Pandey, D. P., Singh, J. P., Roy, R., Singh, V. P., and Pandey, V. B. (1996) Constituents of Heliotropiumindicum, *Oriental Journal of Chemistry*. 12: 321-322.
- Pianowski, L. F., Calixto, J. B., and Chaves, C. P. (2011) *Pharmaceutical Oral Product Obtained from Parts of Heliotropium Plants*. Patent no. WO2011086422A1
- Quisumbing, E. (1951) Medicinal plants of the Philippines, Tech Bull, vol. 16, Pp. 126.
- Rahman, M. A., Mia, M., and Shahid, I. (2011) Pharmacological and phytochemical screen activities of roots of *Heliotropium indicum* Linn. *Pharmacology OnLine*. 1(1): 185–192
- Rao, P. R., Nammi, S., and Raju, A. D. V. (2002) Studies on the antimicrobial activity of *Heliotropiumindicum*Linn. *Journal of Natural Remedies*. 2(2): 195–198.
- Reddy, J. S., Rao, P. R., and Reddy, M. S. (2002) Wound healing effects of *Heliotropiumindicum*, *Plumbagozeylanicum* and *Acalyphaindica*in rats. *Journal of Ethnopharmacology*. 79(2): 249–251
- Sarkar, B. (2016). Ethnic practices and human welfare in India: An attempt for controlling fertility. *International Journal of Experimental Research and Review*. 2: 28-31. doi: https://doi.org/1 Bhattacharjee 0.52756/ijerr.2016.v2.006
- Sarkar, B. (2017). Traditional use of medicinal plants and its biodiversity in India. *International Journal of Experimental Research and Review*. 10: 23-26.
- Sanyal, R., Bala, S. and Mazumdar, A. (2016). Indigenous knowledge of Ethnic community on usage of Satavari (*Asparagus racemosus* Willd) and its preliminary screening.

International Journal of Experimental Research and Review. 7: 62-68.

- Sanyal, R., Mallick, S., and Mazumder, A. (2018). Indigenous Knowledge of Ethnic Community on Usage of Kripa (*Lumnitzera racemosa*) and its preliminary screening. *International Journal of Experimental Research and Review*. 15: 44-50. doi: https://doi.org/10.52756/ijerr.2018.v15.007
- Savadi, R. V., Alagawadi, K. R. and Darade, S. S. (2009) Antifertility activity of ethanolic extract and its n-hexane and benzene fractions of *Heliotropiumindicum*leaves on albino rats. *Journal of Pharmacy Research*. 2 (5): 927–930.
- Shahnaj, S., Asha, U., Mim, T., Rumi, N.S.H., Akter, S., Ghose, S.R., Akter, S., Islam, M.T., Das, P.R., and Rahmatullah, M.(2015). A survey on the ethnomedicinal practices of a folk medicinal practitioner in Manikganj district, Bangladesh. *Journal of Chemical and Pharmaceutical Research*. 7 (8):690–696
- Shalini, S., Kaza, R., and Shaik F. (2010) Study on the anti-inflammatory activity of *Heliotropiumindicum. Journal of Innovative Trends in Pharmaceutical Sciences.* 1(1): 43.
- Souza, J. S. N., Machado, L. L., Pessoa O. D. L., Filho, R. B., Overk, C.R., Yao, P., Cordell, G.A., and Lemos, T.L.G. (2005) Pyrrolizidine alkaloids from *Heliotropiumindicum*. *Journal of the Brazilian Chemical Society*. 16(6B): 1410–1414.
- Srinivas, K., Rao, M. E. B., and Rao, S. (2000) Anti-inflammatory activity of *Heliotropiumindicum* Linn and *Leucasaspera* Spreng. in albino rats. *Indian Journal of Pharmacology*. 32(1):37-38
- Suroowan, S., Pynee, K. B., and Mahomoodally, M. F., (2019) A comprehensive review of ethnopharmacologically important medicinal plant species from Mauritius. *South African Journal of Botany*. 122: 189–213.
- Togola, A., Diallo, D., Demb'el'e, S., Barsett, H., and Paulsen, B. S. (2005) Ethnopharmacological survey of different uses of seven medicinal plants from Mali, (West Africa) in the regions Doila, Kolokani and Siby. *Journal of Ethnobiology and Ethnomedicine*. 1: 7.
- Veda, V. T., Sasi, K. S., Asokan, B. R., Sengottuvelu, S., and Jaikumar, S. (2016) Anticataract activity of ethanolic extract of *Heliotropiumindicum*leaves on galactose induced cataract in rats. *International Journal of Pharmacology & Toxicology*. 5: 18–20

Wiart, C. (2006) Medicinal Plants of the Asia-Pacific, CRC Press, Boca Raton, FL, USA.

Williaman J. J., and Schubert, B. G. (1962) Alkaloid-bearing plants and their contained alkaloids. Technical Bulletin No. 1234. *Journal of Pharmaceutical Sciences*. 51(3): 296.

HOW TO CITE

Rupam Mandal (2022). Heliotropium indicum L: An Ethnomedicinally Important plant of India. © International Academic Publishing House (IAPH), B. Sarkar (eds.), *The Basic Handbook of Indian Ethnobotany and Traditional Medicine*,Vol. 1, pp. 36-45. ISBN: 978-81-957954-1-3. DOI: https://doi.org/10.52756/bhietm.2022.e01.003